

ON THE COMPARISON OF POSITIVE ELEMENTS OF A C*-ALGEBRA BY LOWER SEMICONTINUOUS TRACES

LEONEL ROBERT

ABSTRACT. It is shown in this paper that two positive elements of a C*-algebra agree on all lower semicontinuous traces if and only if they are equivalent in the sense of Cuntz and Pedersen. A similar result is also obtained in the more general case where the two elements are comparable by their values on the lower semicontinuous traces. This result is used to give a characterization of the functions on the cone of lower semicontinuous traces of a stable C*-algebra that arise from positive elements of the algebra.

1. INTRODUCTION

In [1], Cuntz and Pedersen considered the problem of comparing positive elements of a C*-algebra by their values on the lower semicontinuous traces on the algebra. They defined an equivalence relation on the positive elements—the Cuntz-Pedersen relation—and showed, among other results, that if the C*-algebra is simple, then two positive elements are Cuntz-Pedersen equivalent if and only if they agree on all the lower semicontinuous traces. The question whether this was true for an arbitrary C*-algebra was left unsettled in their paper, and is answered affirmatively in Theorem 1 below.

Let A be C*-algebra. Recall that a trace on A is a map $\tau: A^+ \rightarrow [0, \infty]$ that is additive, homogeneous, and satisfies the identity $\tau(xx^*) = \tau(x^*x)$. We will be mostly interested in the lower semicontinuous traces on A , the set of which we shall denote by $T(A)$.

For $a, b \in A^+$ let us say that a is Cuntz-Pedersen equivalent to b , denoted by $a \sim b$, if $a = \sum_{i=1}^{\infty} x_i x_i^*$ and $b = \sum_{i=1}^{\infty} x_i^* x_i$ for some $x_i \in A$. Let us say that a is Cuntz-Pedersen smaller than b , and denote this by $a \preceq b$, if $a \sim a' \leq b$ for some $a' \in A^+$. It was shown in [1] that the relations $\sim$ and $\preceq$ are transitive (so $\sim$ is an equivalence relation).

Theorem 1. *Let a and b be positive elements of A . The following propositions are true.*

- (i) $\tau(a) \leq \tau(b)$ for all $\tau \in T(A)$ if and only if for every $\epsilon > 0$ there is $\delta > 0$ such that $(a - \epsilon)_+ \preceq (b - \delta)_+$.
- (ii) $\tau(a) = \tau(b)$ for all $\tau \in T(A)$ if and only if $a \sim b$.

Remark. We shall see in Section 3 an example where $\tau(a) \leq \tau(b)$ for all $\tau \in T(A)$, but it is not true that $a \preceq b$.

2. PROOF OF THEOREM 1

The proof of Theorem 1 is preceded by a number of preliminary definitions and results.

Let $a, b \in A^+$. Let us write $a \preceq_{CP} b$ if for every $\epsilon > 0$ there is $\delta > 0$ such that $(a - \epsilon)_+ \preceq (b - \delta)_+$. Since $\preceq$ is transitive, the relation $\preceq_{CP}$ is clearly transitive as well. Let us write $a \sim_{CP} b$ if $a \preceq_{CP} b$ and $b \preceq_{CP} a$. This defines an equivalence relation in A^+ . (It will be shown in the proof of Theorem 1 (ii) that $\sim_{CP}$ is the same as $\sim$.)

Proposition 1. *Let $a, b, c, d \in A^+$. The following propositions are true.*

- (i) *If $a \preceq_{CP} b$ and $c \preceq_{CP} d$ then $a + c \preceq_{CP} b + d$.*
- (ii) *If $a \preceq_{CP} b$ then $\alpha a \preceq_{CP} \alpha b$ for all $\alpha \in \mathbb{R}^+$.*
- (iii) *If $a \preceq b$ then $a \preceq_{CP} b$.*

Proof. It was shown in [2, Proposition 2.3] that for all $a, b \in A^+$ and $\epsilon > 0$ there is $\delta > 0$ such that

- (1) $(a - \epsilon)_+ + (b - \epsilon)_+ \preceq (a + b - \delta)_+,$
- (2) $(a + b - \epsilon)_+ \preceq (a - \delta)_+ + (b - \delta)_+.$

These inequalities imply (i).

(ii) This is clear.

(iii) Suppose that $a = \sum_{i=1}^{\infty} x_i x_i^*$ and $\sum_{i=1}^{\infty} x_i^* x_i \leq b$. Let $\epsilon > 0$. By the lemma of Kirchberg and Rørdam [3, Lemma 2.2] (see also the remark after [2, Lemma 2.2]), there are $n \in \mathbb{N}$ and $\epsilon_1 > 0$ such that $(a - \epsilon)_+ \preceq (\sum_{i=1}^n x_i x_i^* - \epsilon_1)_+$. We have

$$\begin{aligned} (a - \epsilon)_+ &\preceq \left(\sum_{i=1}^n x_i x_i^* - \epsilon_1 \right)_+ \preceq \sum_{i=1}^n (x_i x_i^* - \epsilon_2)_+ \sim \sum_{i=1}^n (x_i^* x_i - \epsilon_2)_+ \\ &\preceq \left(\sum_{i=1}^n x_i^* x_i - \epsilon_3 \right)_+ \preceq (b - \epsilon_4)_+. \end{aligned}$$

In the above chain of inequalities we have applied (1), (2) and that $(xx^* - \epsilon)_+ \sim (x^*x - \epsilon)_+$ for all $\epsilon > 0$ (see [2, Proposition 2.3]). $\square$

Let us denote by A_{CP} the quotient $A^+ / \sim_{CP}$. We consider A_{CP} ordered by the order $\langle a \rangle \leq \langle b \rangle$ if $a \preceq_{CP} b$, where $\langle a \rangle$ and $\langle b \rangle$ denote the equivalence classes of the positive elements a and b . We also endow A_{CP} with the addition operation $\langle a \rangle + \langle b \rangle := \langle a + b \rangle$. The order of A_{CP} is compatible with the addition operation, i.e., $\langle a \rangle \leq \langle a \rangle + \langle b \rangle$. Thus, A_{CP} is an ordered semigroup with 0.

In order to prove Theorem 1 (i) we will apply the following proposition to the ordered semigroup A_{CP} .

Proposition 2. ([4, Proposition 3.2]) *Let S be an ordered semigroup with 0 and with the property that if $(k+1)x \leq ky$ for some $x, y \in S$ and $k \in \mathbb{N}$, then $x \leq y$ (i.e., S is almost unperforated). The following implication holds in S :*

If $x \leq My$ for some $M \in \mathbb{N}$, and $\lambda(x) < \lambda(y)$ for every $\lambda: S \rightarrow [0, \infty]$ that is additive, order-preserving, and satisfies $\lambda(y) = 1$, then $x \leq y$.

Notice that the semigroup A_{CP} satisfies the hypotheses of the preceding proposition. In fact, in A_{CP} we have that $k\langle x \rangle \leq k\langle y \rangle$ implies $\langle x \rangle \leq \langle y \rangle$, because we can multiply the elements of A_{CP} by positive real scalars.

Notice also that for every additive map $\lambda: A_{CP} \rightarrow [0, \infty]$, the map on A^+ defined by $\tau_\lambda(a) := \lambda(\langle a \rangle)$ is a trace, because it is additive and satisfies the trace identity (homogeneity holds automatically for any additive map with values in $[0, \infty]$). This trace may not be lower semicontinuous. To Theorem 1 (i) we will then need the following lemma.

Lemma 1. ([2, Lemma 3.1]) *Let $\tau: A^+ \rightarrow [0, \infty]$ be a trace on the C^* -algebra A . Then $\tilde{\tau}(a) = \sup_{\epsilon > 0} \tau((a - \epsilon)_+)$ is a lower semicontinuous trace.*

Proof of Theorem 1 (i). It is clear that if $a \preceq_{CP} b$ then $\tau(a) \leq \tau(b)$ for every $\tau \in T(A)$.

Let us assume that $\tau(a) \leq \tau(b)$ for all $\tau \in T(A)$. For every closed two-sided ideal I of A , the map defined by $\tau_I(x) = 0$ if $x \in I^+$ and $\tau_I(x) = \infty$ otherwise, is a lower semicontinuous trace on A . Since $\tau_I(a) \leq \tau_I(b)$ for any such trace it follows that $\text{Ideal}(a) \subseteq \text{Ideal}(b)$. Let $\epsilon > 0$. We have $(a - \epsilon)_+ \in \text{Ped}(\text{Ideal}(b))^+$; hence $(a - \epsilon)_+ = \sum_{i=1}^m y_i b y_i^*$ for some $y_i \in A$. This implies that $(a - \epsilon)_+ \preceq M b$ for some $M \in \mathbb{N}$, and so $\langle (a - \epsilon)_+ \rangle \leq M \langle b \rangle$. Let us show that we also have $\lambda(\langle (a - \epsilon)_+ \rangle) < \lambda(\langle b \rangle)$ for any additive, order-preserving map $\lambda: A_{CP} \rightarrow [0, \infty]$ such that $\lambda(\langle b \rangle) = 1$. By Proposition 2, this will imply that $\langle (a - \epsilon)_+ \rangle \leq \langle b \rangle$, and since ϵ is arbitrary, we will have $\langle a \rangle \leq \langle b \rangle$, as desired.

Let $\lambda: A_{CP} \rightarrow [0, \infty]$ be additive, order-preserving, and such that $\lambda(\langle b \rangle) = 1$. Let τ_λ be the trace associated to λ and $\tilde{\tau}_\lambda$ its lower semicontinuous regularization defined as in Lemma 1. Notice that for every $\delta > 0$ and $c \in A^+$ we have $\tau_\lambda((c - \delta)_+) \leq \tilde{\tau}_\lambda(c) \leq \tau_\lambda(c)$.

Case 1. Assume that $\tilde{\tau}_\lambda(a) \neq 0$. Then

$$\lambda(\langle (a - \epsilon)_+ \rangle) = \tau_\lambda((a - \epsilon)_+) \leq \tilde{\tau}_\lambda((a - \epsilon/2)_+) < \tilde{\tau}_\lambda(a) \leq \tilde{\tau}_\lambda(b) \leq \tau_\lambda(b) = \lambda(\langle b \rangle).$$

Case 2. Assume that $\tilde{\tau}_\lambda(a) = 0$. Then

$$\lambda(\langle (a - \epsilon)_+ \rangle) = \tau_\lambda((a - \epsilon)_+) = 0 < 1 = \lambda(\langle b \rangle). \quad \square$$

We now turn to the proof of Theorem 1 (ii).

Lemma 2. *Let $a, b, c \in A^+$. The following propositions are true.*

(i) *If $a + c \preceq b + c$ and $c \preceq M b$ for some $M \in \mathbb{R}^+$, then $a \preceq (1 + \delta)b$ for all $\delta > 0$.*

(ii) *If $a + c \sim b + c$ and $c \preceq M a, M b$ for some $M \in \mathbb{R}^+$, then $a \sim b$.*

Proof. (i) Adding a on both sides of $a + c \preceq b + c$ we obtain that $2a + c \preceq 2b + c$, and by induction $ka + c \preceq kb + c$ for all k . Dividing the last inequality by k we see that we may assume that $c \preceq b$. Changing c to c' such that $c \sim c' \leq b$ we may further assume that $c \leq b$. We have $a + b = a + c + (b - c) \preceq 2b$. By applying induction we obtain that $ka + b \preceq (k + 1)b$ for all k . This proves (i).

(ii) As in the proof of (i) we may assume that $c \leq b$. We have $a + b \sim a + c + b - c \sim 2b$. In the same way we obtain that $a + b \sim 2a$. Therefore, $a \sim b$. $\square$

Proof of Theorem 1 (ii). By Theorem 1 (i) we know that $a \preceq_{CP} b$ and $b \preceq_{CP} a$. Thus, there are strictly decreasing sequences of positive numbers ϵ_n and μ_n such that $(a - \epsilon_n)_+ \preceq (b - \mu_n)_+ \preceq (a - \epsilon_{n+1})_+$ for all $n \in \mathbb{N}$, and $\epsilon_n, \mu_n \rightarrow 0$. Let us write $a_n = (a - \epsilon_n)_+$ and $b_n = (b - \mu_n)_+$. The proof now follows the same arguments as in [1, Lemma 7.4], although with some modifications.

Since $a_n \preceq b_n$ there is a'_n such that $a_n \sim a'_n \leq b_n$. We have $(b_n - a'_n) + a'_n \preceq (a_{n+1} - a_n) + a_n$. We also have $a_n \leq M(a_{n+1} - a_n)$. (This is verified by referring back to the definition of a_n in terms of a .) By Lemma 2 (i) we have that $b_n - a'_n \preceq (1 + \delta)(a_{n+1} - a_n)$ for all $\delta > 0$. For δ small enough we have that $b_n - a'_n \preceq (1 + \delta)(a_{n+1} - a_n) \leq a_{n+2} - a_n$. Let z be such that $b_n - a'_n \sim z \leq a_{n+2} - a_n$ and let $b'_n = a_n + z$. The sequence (b'_n) satisfies that $b'_n \sim b_n$ and $a_n \leq b'_n \leq a_{n+2}$.

We have $(b'_{n+3} - b'_n) + b'_n \sim (b_{n+3} - b_n) + b_n$. Let us see that we can apply Lemma 2 (ii) to conclude that $(b'_{n+3} - b'_n) \sim (b_{n+3} - b_n)$. On one hand, $b_n \leq M(b_{n+3} - b_n)$ for some $M > 0$, by the expression of these elements in terms of the functional calculus of b . On the other hand, $b'_n \leq a_{n+2} \leq M'(a_{n+3} - a_{n+2}) \leq M'(b'_{n+3} - b'_n)$ for some $M' > 0$. Hence, Lemma 2 (ii) can be applied. We have

$$a = b'_0 + \sum_{k=1}^{\infty} (b'_{3(k+1)} - b'_{3k}) \sim b_0 + \sum_{k=1}^{\infty} (b_{3(k+1)} - b_{3k}) = b. \quad \square$$

3. FURTHER REMARKS

Let us first show that Theorem 1 (i) cannot be strengthened to obtain $\tau(a) \leq \tau(b)$ for all τ implies $a \preceq b$.

Example. Let A be a simple unital C^* -algebra with exactly two extreme tracial states τ_1 and τ_2 . Such an algebra may be found, say, among the AF C^* -algebras (e.g., [1, Example 6.10]). Let $b \in A^+$ be such that $\tau_1(b) < \tau_2(b)$ and set $\tau_1(b)1 = a$. Then $\tau_1(a) = \tau_1(b)$ and $\tau_2(a) < \tau_2(b)$, and so for every bounded trace τ , $\tau(a) \leq \tau(b)$. This also holds for the unique unbounded trace with value ∞ on all points except 0. But we cannot have $a \preceq b$, because in that case $a + c \sim b$ for some $c \in A^+$, and so $\tau_1(c) = 0$ and $\tau_2(c) > 0$, which is impossible by the simplicity of A .

Let us consider the following topology on $T(A)$: the net of lower semicontinuous traces (τ_i) converges to τ if

$$\limsup \tau_i((a - \epsilon)_+) \leq \tau(a) \leq \liminf \tau_i(a)$$

for all $a \in A^+$ and $\epsilon > 0$. It was shown in [2] that $T(A)$ is a compact Hausdorff space and that the functor $T(\cdot)$ is a continuous contravariant functor from the category of C^* -algebras to the category of topological spaces.

For every positive element $a \in A^+$ the function $\tilde{a}: T(A) \rightarrow [0, \infty]$, defined by $\tilde{a}(\tau) = \tau(a)$, is linear and lower semicontinuous. Consider the following question: Which linear and lower semicontinuous functions on $T(A)$ arise from positive elements of A in the form $\tilde{a}$? In view of Theorem 1, this question is asking for a description of the ordered semigroup A_{CP} as a semigroup of

functions on $T(A)$. Theorem 2 below gives an answer to this question assuming that A is stable.

Let us recall the definition of the non-cancellative cone $S(T(A))$ given in [2]. Let $LSC(T(A))$ denote the linear, lower semicontinuous, functions on $T(A)$ with values in $[0, \infty]$. Then $S(T(A))$ is composed of the functions $f \in LSC(T(A))$ that satisfy the following condition: there is an increasing sequence (h_n) , $h_n \in LSC(T(A))$, such that $f = \sup h_n$ and h_n is continuous on each point where h_{n+1} is finite.

For every $a \in A^+$ we have $\tilde{a} \in S(T(A))$ (see [2, Proposition 5.1]). Moreover, by [2, Theorem 5.9], $S(T(A))$ is the set of functions $f \in LSC(T(A))$ for which there is an increasing sequence of functions $\tilde{a}_n$ coming from positive elements $a_n \in A^+$ and such that $f = \sup \tilde{a}_n$. We can now prove the following theorem.

Theorem 2. *Let A be a stable C^* -algebra. Then $S(T(A))$ is the set of functions of the form $\tilde{a}$ for some $a \in A^+$.*

Proof. Let f be in $S(T(A))$ and let $(\tilde{a}_n)$, with $a_n \in A^+$, be an increasing sequence of functions with supremum f . By Theorem 1 (i), we have that $a_n \preceq_{CP} a_{n+1}$ for all n . We may replace the positive elements a_n by $a'_n = (a_n - \epsilon_n)_+$, with $\epsilon_n > 0$ small enough, so that $a'_n \preceq a'_{n+1}$ and $f = \sup \tilde{a}'_n$. Let $c_n \in A^+$ be such that $a'_n + c_n \sim a'_{n+1}$. Using the stability of A we may find mutually orthogonal elements c''_n such that $c''_n \sim c'_n$. Furthermore, again by the stability of A , we may assume that the c''_n s have sufficiently small norm such that the series $a = \sum_{n=1}^{\infty} c''_n$ converges. We then have $f = \tilde{a}$. $\square$

Question 1. Is $S(T(A)) = LSC(T(A))$?

Question 2. In [2] a cone dual to the 2-quasitraces of A is defined which is analogous to the cone $S(T(A))$. Is the analog of Theorem 2 for 2-quasitraces true?

Acknowledgments. I am grateful to George A. Elliott for pointing out that Theorem 2 was in all likelihood true.

REFERENCES

- [1] J. Cuntz and G. K. Pedersen, *Equivalence and traces on C^* -algebras*, J. Funct. Anal. **33** (1979), no. 2, 135–164.
- [2] G. A. Elliott, L. Robert, and L. Santiago, *On the cone of lower semicontinuous traces on a C^* -algebra*, arXiv:0805.3122v1 [math. OA] (2008).
- [3] E. Kirchberg and M. Rørdam, *Infinite non-simple C^* -algebras: absorbing the Cuntz algebras $\mathcal{O}_\infty$* , Adv. Math. **167** (2002), no. 2, 195–264.
- [4] M. Rørdam, *The stable and the real rank of $\mathcal{Z}$ -absorbing C^* -algebras*, Internat. J. Math. **15** (2004), no. 10, 1065–1084.

LEONEL ROBERT, FIELDS INSTITUTE, TORONTO, CANADA, M5T 3J1
E-mail address: lrobert@fields.utoronto.ca